

Signature:

Darlinghurst NSW 2010
Tel: 02 9360 4811 • fax: 02 9360 9919

Ground floor, 67 Burton Street

Web: http://sydneyentclinic.com

Patient	Appointment
---------	-------------

Test(s) required (please note that challenge tests must be performed separately) For revision airway surgery, pre/post rhinoplasty, □ nasal airway medicolegal documentation, OSA investigations, pre/post CPAP use (decongestion test with acoustic **ASSESSMENT** rhinometry, rhinomanometry & nasendoscopy). Inhalant allergy screen suitable for northern NSW to ANTIGEN SKIN PRICK TEST VIC, with assessment of house dust mites, animal epithelia, moulds, weeds, tree pollens, temperate and tropical grass pollens. IMMUNOTHERAPY Discussion of suitability for sublingual immunotherapy. When clinical suspicion exists but RAST and DIRECT NASAL ANTIGEN epicutaneous testina non-confirmatory. are CHALLENGE Suspected antigen must be specified in comments. Three evoked olfactory responses: odour threshold, OLFACTORY ASSESSMENT identification and discrimination, with validated patient outcome questionnaires. Discussion of suitability for smell re-training and SMELL RE-TRAINING rehabilitation. Identification and threshold tests assessing sweet, TASTE ASSESSMENT sour, salty and bitter taste. L LYSINE ASPIRIN Identification of aspirin sensitive airways disease. Please bring your asthma medication. **CHALLENGE** ADDITIONAL RHINOLOGIC REQUIRES SEPARATE APPOINTMENT IF REQUESTED IN CONJUNCTION WITH TESTING. **OPINION WITH PROF HARVEY** Paranasal sinus and airway scan. Non-contrast. Bone LOW DOSE CONICAL BEAM windows. Open frame scanner. 0.17mSv total dose CT SCAN (SINUSES) (1mSv+ for traditional CT). Referral date: Referring doctor:

Read patient information over page before your appointment.

Provider number:

PATIENT INFORMATION

TEST	APPROX DURATION	NOTES
Airway assessment	1 hour	Avoid nasal/oral decongestants (eg Drixine, Otrivin, Nyal etc) or any tablets containing pseudoephedrine for 7 days prior .
Challenge test	2 hours, plus monitoring	Avoid nasal/oral decongestants and pseudoephedrine , as above. Avoid antihistamines (eg Claratyne, Telfast, Zyrtec) for <u>7 days prior</u> .
Skin prick testing (allergy)	45 minutes	Avoid antihistamines, as above. Some sleeping medications, blood pressure/heart medications, and antidepressants may interfere with testing. Please contact us prior to testing.
Olfactory assessment	60 minutes	Do not eat for at least 30mins prior to test.
Taste assessment	30 minutes	Do not eat for at least 30mins prior to test.

How to find us:

PARKING

- On-street parking is limited
- St. Vincent's Public Hospital Car Park, enter via Victoria Street (fees apply) PUBLIC TRANSPORT



